




U R B A N P E S T C O N T R O L

TERMIDOR® 2,5% EC

For fast and effective treatment of
subterranean termites

Solve termite problems within 3 months by using TERMIDOR®, an emulsifiable concentrate that kills by residual contact or ingestion!

TERMIDOR® contains fipronil for the control of wood destroying termites damaging buildings or structures, or subterranean termites infesting earthen dam walls.

In a class of its own

After nine years of field tests, TERMIDOR® produced 100 % control of termites, with no signs of re-infestation at any of the study sites involved. Colonies managed in 3 months or less. Termidor produces effective results up to 6 times faster than bait systems!

The ultimate solution

TERMIDOR® is unique, in that it is a new technology liquid that is undetectable to termites. This invincible killer – invisible, odourless and tasteless kills both by residual contact and ingestion. As termites go about their normal routines they contact and ingest TERMIDOR®, thereby transferring it to other termites that they come into contact with throughout the colony. TERMIDOR® has been designed to be slow acting, thereby ensuring that termites have ample time to spread it throughout their entire population. This “carrier” system is known as the “Transfer Effect,” and results in 100 % control. Responsible, low-dose treatment that builds client confidence!

Superior to traditional liquid products and baits

Traditional liquid termiticides are repellents, which termites can not only detect, but also avoid. TERMIDOR®, on the other hand, is undetectable, therefore termites neither avoid it nor

try to circumvent it, and forage freely in the treated zone. As a result, the “Transfer Effect,” begins and in turn, the elimination of the colony. Many bait systems work by means of “recruitment,” and as a result can take from many months to several years to control a colony. Termites first have to discover the bait station and recruit sufficient members to it, before a termite control product is added to the station. Furthermore, there is no guarantee that termites will even find the bait station. TERMIDOR®, on the other hand, starts working immediately and provides control within 3 months.

Choose TERMIDOR® - America's No. 1 termiticide with over 1 million homes treated since 2000!

Locations serviced by Termidor


Foundation walls and trenches, soil surfaces, under floor slabs, suspended floors, pile and beam construction, unexposed soil surfaces, earthen dam walls, termite mounds and colonies all respond very positively to TERMIDOR® treatment.

Peace of mind

Each time the pest control operator tackles termites, he places his reputation on the line – Choose effective TERMIDOR® and reduce termite-related callbacks, thereby increasing profitability, with minimum fuss, yet maximum results within 3 months!



TERMIDOR®
TERMITE PROTECTION™





TERMIDOR®
TERMITE PROTECTION™

Technical Information

FOR PROFESSIONAL USE ONLY: Only for use by registered pest control operators.

Formulation and use indications: An emulsifiable concentrate residual contact and stomach insecticide for the control of wood destroying subterranean termites damaging buildings and structures.

Active Ingredient: Fipronil (phenyl pyrazole) 25 g/l

Toxicology: WHO classification – Class II – slightly hazardous. Fipronil binds more tightly to insect receptors than to vertebrate ones, so it is more toxic to insects than vertebrates.

Application:


General directions: TERMIDOR® must be applied according to SABS Code of Practice 0124:1995 as amended. All applications must be made with suitable equipment, designed for applications for termite control, which is in good working order and correctly calibrated to give the desired coverage.

Pre-construction preventive application

Treatment of existing colonies: All existing active termite colonies situated within 50 m of the structures perimeter should be treated with TERMIDOR® 2,5% EC (see below). The treatment of surrounding nests will assist in preventing attack on structures subsequently erected on the site.

All cellulose containing material (wood, paper, cardboard etc.) should be removed from the site before treatment. Do not apply TERMIDOR® 2,5% EC to excessively wet soil. If more than 15 mm rain falls within 24 hours after application, the application should be repeated unless the treated soil was adequately protected.

In all cases the treated soil surface should not be disturbed after treatment, as this may damage the treated barrier, allowing termites access to the structure through such areas.



Location	Dosage / 100 / water	Method of application
Foundation and service trenches and foundation walls	1 /	Treat service trenches and inner foundation walls at the rate of 5 / mixture per linear meter.
Soil surface under floor slabs	1 /	Apply 5 / mixture per 1 m ² on the soil surface where the floor will be cast.
Unexposed soil surfaces	1 /	Apply 5 / of the mixture per 1 m ² on the soil surface where paving, entrance steps, verandas etc. will be connected to the structure.

Post-construction remedial application

With remedial treatments it is important that the application is carried out in such a way that the TERMIDOR® 2,5% EC is distributed in a continuous layer in the soil under existing structures.

Location	Dosage / 100 / water	Method of application
Perimeter treatment	1 /	Dig a 30 cm wide trench around the exterior perimeter of the foundations and as deep as the top of the footing. Apply 5 / mixture per linear metre of the trench.
Solid floors	1 /	Drill holes approximately 1 m apart through the floor around the inside perimeter of each room, and around the outside perimeter of the structure. Apply 5 / mixture per hole. Holes should be suitably sealed after application.
Suspended floors	1 /	A 30 cm wide trench should be dug around the inside of the foundation walls as deep as the top of the footing and treated with 5 / mixture per linear metre. Treat the surface of the crawl space and the enclosing walls at a rate of 5 litres mixture per m ² .

Treatment of existing termite mounds / colonies

TERMIDOR® 2,5% EC can be used to control existing active colonies by treating the nests.

Application rate: Mix 50 m TERMIDOR® 2,5% EC in 5 / water (1 / in 100 / water) and apply 5 / mixture to the termite nest as follows:

- **Termite mounds:** Remove the top of the mound with a spade or small axe and pour 5 / of the mixture into the termite nest.
- **Termite mounds with "chimneys" (e.g. *Odontotermes* spp.):** Pour 5 / mixture into as many chimneys as can be found. Divide the 5 / mixture evenly between the various chimneys.

Always read and follow label directions

TERMIDOR® 2,5% EC: Reg.No. L6616 (Act 36/1947) contains fipronil (phenyl pyrazole) 25g/l. Harmful

Registered by : BASF SA (Pty) Ltd. Co. Reg. No. 1966/10235/07.
P.O. Box 2801, Halfway House, 1685. Tel. 011 203-2400
Manufactured & formulated by BASF Agro BV, Arnheim (NL), Waedenswil Branch,
Steinacherstrasse 101, CH-8820 Waedenswil, Switzerland.

® is the registered trademark of BASF AG, Germany.